

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 October 2002 (24.10.2002)

PCT

(10) International Publication Number
WO 02/083677 A1

(51) International Patent Classification⁷: **C07D 487/04**,
A01N 43/90 // (C07D 487/04, 249:00, 239:00)

(21) International Application Number: PCT/EP02/03830

(22) International Filing Date: 6 April 2002 (06.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01109010.7 11 April 2001 (11.04.2001) EP

(71) Applicant (for all designated States except US): **BASF AKTIENGESELLSCHAFT** [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **TORMO I BLASCO, Jordi** [ES/DE]; Mühlweg 47, 67117 Limburgerhof (DE). **SAUTER, Hubert** [DE/DE]; Neckarpromenade 20, 68167 Mannheim (DE). **MÜLLER, Bernd** [DE/DE]; Jean-Ganss-Strasse 21, 67227 Frankenthal (DE). **GEWEHR, Markus** [DE/DE]; Goethestrasse 21, 56288 Kastellaun (DE). **GRAMMENOS, Wassilios** [GR/DE]; Borsigstrasse 5, 67063 Ludwigshafen (DE). **GROTE, Thomas** [DE/DE]; Im Hoehnhausen 18, 67157 Wachenheim (DE). **GYPSE, Andreas** [DE/DE]; B 4,4, 68159 Mannheim (DE). **RHEINHEIMER, Joachim** [DE/DE]; Merziger Strasse 24, 67063 Ludwigshafen (DE). **ROSE, Ingo** [DE/DE]; C 2, 19, 68159 Mannheim (DE). **SCHÄFER, Peter** [DE/DE]; Römerstrasse 1, 67308 Ottersheim (DE). **SCHIEWECK, Frank** [DE/DE]; Lindenweg

4, 67258 Hessheim (DE). **AMMERMAN, Eberhard** [DE/DE]; Von-Gagern-Strasse 2, 64646 Heppenheim (DE). **STRATHMANN, Siegfried** [DE/DE]; Donnersbergstrasse 9, 67117 Limburgerhof (DE). **LORENZ, Gisela** [DE/DE]; Erlengweg 13, 67434 Hambach (DE). **STIERL, Reinhard** [DE/DE]; Ginsterstrasse 17, 67112 Mutterstadt (DE).

(74) Common Representative: **BASF AKTIENGESELLSCHAFT**; 67056 Ludwigshafen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 6-(2-CHLORO-6-FLUORO-PHENYL)-TRIAZOLOPYRIMIDINES



(I)


(57) Abstract: 6-(2-Chloro-6-fluoro-phenyl)-triazolopyrimidines of formula (I), in which R¹ and R² independently denote hydrogen or alkyl, alkenyl, alkynyl, alkadienyl, or haloalkyl, cycloalkyl, bicycloalkyl, phenyl, naphthyl, or 5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or 5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or where R¹ and R² radicals may be unsubstituted or partially or fully halogenated or may be substituted as defined in the description, R¹ and R² together with the interjacent nitrogen atom represent a 5- or

6-membered heterocyclic ring, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, which may be substituted; X is cyano, alkoxy, haloalkoxy or alkenyloxy; processes for their preparation, compositions containing them and to their use for combating phytopathogenic fungi.

6-(2-Chloro-6-fluoro-phenyl)-triazolopyrimidines

Description

The invention relates to 6-(2-chloro-6-fluoro-phenyl)-triazolopyrimidines of formula I



I

in which

R^1 and R^2 independently denote hydrogen or

C_1 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_2 - C_{10} -alkynyl, C_4 - C_{10} -alkadienyl, or C_1 - C_{10} -haloalkyl,

C_3 - C_8 -cycloalkyl, C_5 - C_{10} -bicycloalkyl, phenyl, naphthyl, or 5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or

5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or

where R^1 and R^2 radicals may be unsubstituted or partially or fully halogenated or may carry one to three groups R^a ,

R^a is halogen, cyano, nitro, hydroxyl, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino, C_2 - C_6 -alkenyl, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyl, C_3 - C_6 -alkynyloxy and C_1 - C_4 -alkylenedioxy, which may be halogenated; or

R^1 and R^2 together with the interjacent nitrogen atom represent a 5- or 6-membered heterocyclic ring, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, which may be substituted by one to three R^a radicals;

X is cyano, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy or C_3 - C_8 -alkenyloxy.

Moreover, the invention relates to processes for their preparation, compositions containing them and to their use for combating phytopathogenic fungi.

- 5 EP-A 071 792 discloses 6-phenyl-7-amino-triazolopyrimidines where the 5-position is substituted by hydrogen or alkyl or aryl groups.

EP-A 550 113 relates to 5-H- and 5-halogen-6-phenyl-7-amino-tri-
10 azolopyrimidines where the 7-amino group is further substituted.

WO-A 98/46607 discloses triazolopyrimidines, which are substituted in the 6-position by a 2,4,6-trifluorophenyl group.

- 15 US 5,994,360 discloses triazolopyrimidines, which are substituted in the 5-position by alkyl groups.

The compounds disclosed in the documents discussed above are said to be active against various phytopathogenic fungi.

20

It is an object of the present invention to provide compounds having improved fungicidal activity.

- We have found that this object is achieved by the compounds defined at the outset. Furthermore, we have found processes for their
25 preparation, compositions comprising them and methods for controlling phytopathogenic fungi using the compounds I.


The compounds of the formula I differ from the compounds known
30 from the abovementioned art in the combination of the 2-chloro-6-fluoro-phenyl group and a specific substitution in 5-position of the triazolopyrimidine system.

- A 4- to 6- membered heterocyclic group may be any heterocyclic
35 group with 4 to 6 ring atoms, interrupted by one or more heteroatoms selected from sulfur, nitrogen, and oxygen, preferably oxygen. A halogen atom suitable denotes a fluorine, chlorine or bromine atom.

- 40 The present invention further provides a process for the preparation of compounds of formula I as defined above which comprises treating a 5-halogen compound of formula II

45

3



in which Y is halogen with a compound of formula III, which is, dependent from the value of X to be introduced to yield formula I compounds, an anorganic cyano salt, an alkoxylate, haloalkoxylate or an alkenyloxylate, respectively, preferably in the presence of a solvent. The cation M in formula III has minor influence; for practical and economical reasons usually ammonium-, tetraalkylammonium- or alkalimetal- and earth metal salts are preferred.

15 Suitable solvents include ethers, such as dioxane, diethyl ether and, especially, tetrahydrofuran, halogenated hydrocarbons such as dichloromethane and aromatic hydrocarbons, for example toluene.

20 The reaction is suitably carried out at a temperature in the range from 0 to 120°C, the preferred reaction temperature being from 10 to 40°C.

It is also preferred that the reaction is carried out in the presence of a base. Suitable bases include tertiary amines, such as triethylamine, and inorganic bases, such as potassium carbonate or sodium carbonate. Alternatively, an excess of the compound of formula III may serve as a base.

30 The reaction mixtures are worked up in a customary manner, for example by mixing with water, phase separation and, if required, chromatographic purification of the crude products. Some of the end products are obtained in the form of colorless or slightly brownish, viscous oils, which are purified or freed from volatile components under reduced pressure and at moderately elevated temperatures. If the end products are obtained as solids, purification can also be carried out by recrystallization or digestion.

Compounds of formula II are known in the art and can be obtained by synthesis routes disclosed in EP-A 550 113 and WO-A 98/46608.

If individual compounds I are not obtainable by the routes described above, they can be prepared by derivatization of other compounds I.

45

In the symbol definitions given in the formulae above, collective terms were used which generally represent the following substituents:

5 - halogen: fluorine, chlorine, bromine and iodine;

- C₁-C₁₀-alkyl and the alkyl moieties of C₁-C₁₀-haloalkyl: saturated, straight-chain or branched hydrocarbon radicals having 1 to 10, especially 1 to 6 carbon atoms, for example C₁-C₄-alkyl
10 as mentioned above or pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-di-methylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl,
15 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl;

- C₁-C₆-haloalkyl and the haloalkyl moieties of C₁-C₆-haloalkoxy:
20 straight-chain or branched alkyl groups having 1 to 6, preferably 1 to 4 carbon atoms (as mentioned above), where the hydrogen atoms in these groups may be partially or fully replaced by halogen atoms as mentioned above, for example C₁-C₂-haloalkyl, such as chloromethyl, bromomethyl, dichloromethyl,
25 trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl,
30 2,2,2-trichloroethyl and pentafluoroethyl;

- C₃-C₁₀-alkenyl: unsaturated, straight-chain or branched hydrocarbon radicals having 3 to 10, especially 3 to 6 carbon atoms and a double bond in any position, for example 1-propenyl,
35 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl and 2-methyl-2-propenyl;

- C₂-C₁₀-alkynyl: straight-chain or branched hydrocarbon radicals
40 having 2 to 10, especially 2 to 4 carbon atoms and a triple bond in any position, for example ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne and 1-methyl-2-propynyl;

- In general terms, the term cycloalkyl, as used herein with respect to a radical or moiety refers to a cycloalkyl group having
45 3 to 8 carbon atoms, preferably 5 to 7 carbon atoms.

- In general terms, the term bicycloalkyl, as used herein with respect to a radical or moiety refers to a bicycloalkyl group having 5 to 10 carbon atoms, preferably 6 to 9 carbon atoms, in particular bicycloheptyl being optionally substituted by one or more halogen atoms, nitro, cyano, alkyl, preferably C₁₋₆ alkyl, alkoxy, preferably C₁₋₆ alkoxy.
- 5-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom:
- 10 5-membered heteroaryl groups which, in addition to carbon atoms, may contain one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom as ring members, for example 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-imidazolyl, 4-imidazolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2-yl and 1,3,4-triazol-2-yl;
- 6-membered heteroaryl, containing one to four nitrogen atoms:
- 25 6-membered heteroaryl groups which, in addition to carbon atoms, may contain one to three or one to four nitrogen atoms as ring members, for example 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 1,3,5-triazin-2-yl and 1,2,4-triazin-3-yl.
- 30 With respect to their intended use, preference is given to triazolopyrimidines of the formula I having the following substituents, where the preference is valid in each case on its own or in combination:
- 35 The particularly preferred embodiments of the intermediates with respect to the variables correspond to those of the radicals X, R¹ and R² of formula I.
- A preferred alkyl moiety is an ethyl or especially a methyl
- 40 group.
- A preferred haloalkyl moiety is the 2,2,2-trifluoroethyl or 1,1,1-trifluoroprop-2-yl group;
- 45 A preferred alkenyl moiety is allyl or especially a 2-methylallyl group.

6

A preferred cycloalkyl moiety is cyclopentyl being optionally substituted by one or more halogen atoms, nitro, cyano, alkyl, preferably C₁-C₆ alkyl, alkoxy, preferably C₁-C₆ alkoxy.

5 A preferred heteroaryl moiety is pyridyl, pyrimidyl, pyrazolyl or thienyl.

Preference is given to compounds of formula I in which any alkyl or haloalkyl part of the groups R¹ or R², which may be straight
10 chained or branched, contains up to 10 carbon atoms, preferably 1 to 9 carbon atoms, more preferably 2 to 6 carbon atoms, any alkenyl or alkynyl part of the substituents R¹ or R² contains up to 10 carbon atoms, preferably 2 to 9 carbon atoms, more preferably 3 to 6 carbon atoms, any cycloalkyl part of the substituents
15 R¹ or R² contains from 3 to 10 carbon atoms, preferably from 3 to 8 carbon atoms, more preferably from 3 to 6 carbon atoms, and any bicycloalkyl part of the substituents R¹ or R² contains from 5 to 9 carbon atoms, preferably from 7 to 9 carbon atoms. Any alkyl, alkenyl or alkynyl group may be linear or branched.

20

Likewise, preference is given to compounds of formula I wherein R¹ is not hydrogen.

Compounds of formula I are preferred in which R¹ represents a
25 straight-chained or branched C₁-C₁₀-alkyl, in particular a branched C₃-C₁₀-alkyl group, a C₃-C₈-cycloalkyl, a C₅-C₉-bicycloalkyl, a C₃-C₈-cycloalkyl-C₁-C₆-alkyl, C₁-C₁₀-alkoxy-C₁-C₆-alkyl, a C₁-C₁₀-haloalkyl or a phenyl group being optionally substituted by one to three halogen atoms or C₁-C₁₀-alkyl or C₁-C₁₀-alkoxy groups.

30

Particular preference is given to compounds I in which R² represents a hydrogen atom, a C₁-C₁₀-alkyl or a C₁-C₁₀-haloalkyl group, in particular a hydrogen atom.

35 Besides, particular preference is given to compounds I in which R² is hydrogen.

Moreover, particular preference is given to compounds I in which R² is methyl.

40

Furthermore, particular preference is given to compounds I in which R² is ethyl.

45

If R¹ denotes a C₁-C₁₀-haloalkyl group, preferably a polyfluorinated alkyl group, in particular a 2,2,2-trifluoroethyl, a 2-(1,1,1-trifluoropropyl) or a 2-(1,1,1-trifluorobutyl) group, R² preferably represents a hydrogen atom.

5

If R¹ denotes an optionally substituted C₃-C₈-cycloalkyl group, preferably a cyclopentyl or cyclohexyl group, R² preferably represents a hydrogen atom or C₁-C₆-alkyl group.

- 10 Moreover, particular preference is given to compounds I in which R¹ and R² together with the interjacent nitrogen atom form an optionally substituted heterocyclic ring, preferably an optionally substituted C₃-C₇-heterocyclic ring, in particular a pyrrolidine, piperidine, tetrahydropyridine, in particular 1,2,3,6-tetrahydro-
15 pyridine or azepane ring which is optionally substituted by one or more C₁-C₁₀-alkyl groups.

- Included in the scope of the present invention are (R) and (S) isomers of compounds of general formula I having a chiral center
20 and the racemates thereof, and salts, N-oxides and acid addition compounds.

- With respect to their use, particular preference is given to the compounds I compiled in the tables below. The groups mentioned in
25 the tables for a substituent are furthermore for their part, independently of the combination in which they are mentioned, a particularly preferred embodiment of the respective substituents.

Table 1

- 30 Compounds of the formula I, in which X is cyano and R¹ and R² correspond to one row in Table A

Table 2

- Compounds of the formula I, in which X is methoxy and R¹ and R²
35 correspond to one row in Table A

Table 3

- Compounds of the formula I, in which X is ethoxy and R¹ and R² correspond to one row in Table A
40

Table 4

- Compounds of the formula I, in which X is n-propoxy and R¹ and R² correspond to one row in Table A

45 Table 5

- Compounds of the formula I, in which X is iso-propoxy and R¹ and R² correspond to one row in Table A

Table 6

Compounds of the formula I, in which X is allyloxy and R¹ and R² correspond to one row in Table A

5 Table 7

Compounds of the formula I, in which X is 3-methylallyloxy and R¹ and R² correspond to one row in Table A

Table 8

10 Compounds of the formula I, in which X is fluoromethoxy and R¹ and R² correspond to one row in Table A


Table 9

15 Compounds of the formula I, in which X is difluoromethoxy and R¹ and R² correspond to one row in Table A

Table 10

20 Compounds of the formula I, in which X is trifluoromethoxy and R¹ and R² correspond to one row in Table A

Table A



I

25

No.	R ¹	R ²
30 A-1	CH ₂ CH ₃	H
A-2	CH ₂ CH ₃	CH ₃
A-3	CH ₂ CH ₃	CH ₂ CH ₃
A-4	CH ₂ CF ₃	H
35 A-5	CH ₂ CF ₃	CH ₃
A-6	CH ₂ CF ₃	CH ₂ CH ₃
A-7	CH ₂ CCl ₃	H
A-8	CH ₂ CCl ₃	CH ₃
A-9	CH ₂ CCl ₃	CH ₂ CH ₃
40 A-10	CH ₂ CH ₂ CH ₃	H
A-11	CH ₂ CH ₂ CH ₃	CH ₃
A-12	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
A-13	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃
45 A-14	CH(CH ₃) ₂	H
A-15	CH(CH ₃) ₂	CH ₃
A-16	CH(CH ₃) ₂	CH ₂ CH ₃

No.	R ¹	R ²
A-17	(±) CH(CH ₃)-CH ₂ CH ₃	H
A-18	(±) CH(CH ₃)-CH ₂ CH ₃	CH ₃
5 A-19	(±) CH(CH ₃)-CH ₂ CH ₃	CH ₂ CH ₃
A-20	(R) CH(CH ₃)-CH ₂ CH ₃	H
A-21	(R) CH(CH ₃)-CH ₂ CH ₃	CH ₃
A-22	(R) CH(CH ₃)-CH ₂ CH ₃	CH ₂ CH ₃
10 A-23	(S) CH(CH ₃)-CH ₂ CH ₃	H
A-24	(S) CH(CH ₃)-CH ₂ CH ₃	CH ₃
A-25	(S) CH(CH ₃)-CH ₂ CH ₃	CH ₂ CH ₃
A-26	(±) CH(CH ₃)-CH(CH ₃) ₂	H
A-27	(±) CH(CH ₃)-CH(CH ₃) ₂	CH ₃
15 A-28	(±) CH(CH ₃)-CH(CH ₃) ₂	CH ₂ CH ₃
A-29	(R) CH(CH ₃)-CH(CH ₃) ₂	H
A-30	(R) CH(CH ₃)-CH(CH ₃) ₂	CH ₃
A-31	(R) CH(CH ₃)-CH(CH ₃) ₂	CH ₂ CH ₃
20 A-32	(S) CH(CH ₃)-CH(CH ₃) ₂	H
A-33	(S) CH(CH ₃)-CH(CH ₃) ₂	CH ₃
A-34	(S) CH(CH ₃)-CH(CH ₃) ₂	CH ₂ CH ₃
A-35	(±) CH(CH ₃)-C(CH ₃) ₃	H
25 A-36	(±) CH(CH ₃)-C(CH ₃) ₃	CH ₃
A-37	(±) CH(CH ₃)-C(CH ₃) ₃	CH ₂ CH ₃
A-38	(R) CH(CH ₃)-C(CH ₃) ₃	H
A-39	(R) CH(CH ₃)-C(CH ₃) ₃	CH ₃
A-40	(R) CH(CH ₃)-C(CH ₃) ₃	CH ₂ CH ₃
30 A-41	(S) CH(CH ₃)-C(CH ₃) ₃	H
A-42	(S) CH(CH ₃)-C(CH ₃) ₃	CH ₃
A-43	(S) CH(CH ₃)-C(CH ₃) ₃	CH ₂ CH ₃
A-44	(±) CH(CH ₃)-CF ₃	H
35 A-45	(±) CH(CH ₃)-CF ₃	CH ₃
A-46	(±) CH(CH ₃)-CF ₃	CH ₂ CH ₃
A-47	(R) CH(CH ₃)-CF ₃	H
A-48	(R) CH(CH ₃)-CF ₃	CH ₃
40 A-49	(R) CH(CH ₃)-CF ₃	CH ₂ CH ₃
A-50	(S) CH(CH ₃)-CF ₃	H
A-51	(S) CH(CH ₃)-CF ₃	CH ₃
A-52	(S) CH(CH ₃)-CF ₃	CH ₂ CH ₃
45 A-53	(±) CH(CH ₃)-CCl ₃	H
A-54	(±) CH(CH ₃)-CCl ₃	CH ₃
A-55	(±) CH(CH ₃)-CCl ₃	CH ₂ CH ₃

10

No.	R ¹	R ²
A-56	(R) CH(CH ₃)-CCl ₃	H
A-57	(R) CH(CH ₃)-CCl ₃	CH ₃
5 A-58	(R) CH(CH ₃)-CCl ₃	CH ₂ CH ₃
A-59	(S) CH(CH ₃)-CCl ₃	H
A-60	(S) CH(CH ₃)-CCl ₃	CH ₃
A-61	(S) CH(CH ₃)-CCl ₃	CH ₂ CH ₃
10 A-62	CH ₂ C(CH ₃)=CH ₂	H
A-63	CH ₂ C(CH ₃)=CH ₂	CH ₃
A-64	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃
A-65	cyclopentyl	H
A-66	cyclopentyl	CH ₃
15 A-67	cyclopentyl	CH ₂ CH ₃
A-68	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -	

The compounds I are suitable as fungicides. They have outstanding activity against a broad spectrum of phytopathogenic fungi, in particular from the classes of the *Ascomycetes*, *Deuteromycetes*, *Phycomycetes* and *Basidiomycetes*. Some of them act systemically, and they can be employed in crop protection as foliar- and soil-acting fungicides.

They are especially important for controlling a large number of fungi on a variety of crop plants such as wheat, rye, barley, oats, rice, maize, grass, bananas, cotton, soya, coffee, sugar cane, grapevines, fruit species, ornamentals and vegetables such as cucumbers, beans, tomatoes, potatoes and cucurbits, and on the seeds of these plants.

Specifically, they are suitable for controlling the following plant diseases:

- Alternaria species on vegetables and fruit,
- 35 Botrytis cinerea (gray mold) on strawberries, vegetables, ornamentals and grapevines,
- Cercospora arachidicola on peanuts,
- Erysiphe cichoracearum and Sphaerotheca fuliginea on cucurbits,
- Blumeria graminis (powdery mildew) on cereals,
- 40 Fusarium and Verticillium species on various plants,
- Helminthosporium species on cereals,
- Mycosphaerella species on bananas and peanuts,
- Phytophthora infestans on potatoes and tomatoes,
- Plasmopara viticola on grapevines,
- 45 Podosphaera leucotricha on apples,
- Pseudocercospora herpotrichoides on wheat and barley,
- Pseudoperonospora species on hops and cucumbers,

11

- Puccinia species on cereals,
Pyricularia oryzae on rice,
Rhizoctonia species on cotton, rice and lawns,
Septoria species in cereals,
5 Uncinula necator on grapevines,
Ustilago species on cereals and sugar cane, and
Venturia species (scab) on apples and pears.

- Moreover, the compounds I are suitable for controlling harmful
10 fungi such as Paecilomyces variotii in the protection of materials (e.g. wood, paper, paint dispersions, fibers and tissues) and in the protection of stored products.

- The compounds I are applied by treating the fungi, or the plants,
15 seeds, materials or the soil to be protected against fungal infection, with a fungicidally active amount of the active ingredients. Application can be effected both before and after infection of the materials, plants or seeds by the fungi.

- 20 In general, the fungicidal compositions comprise from 0.1 to 95, preferably 0.5 to 90, % by weight of active ingredient.

- When used in crop protection, the rates of application are from 0.01 to 2.0 kg of active ingredient per ha, depending on the nature
25 of the effect desired.

- In the treatment of seed, amounts of active ingredient of from 0.001 to 0.1 g, preferably 0.01 to 0.05 g, are generally required per kilogram of seed.

- 30 When used in the protection of materials or stored products, the rate of application of active ingredient depends on the nature of the field of application and on the effect desired. Rates of application conventionally used in the protection of materials are,
35 for example, from 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active ingredient per cubic meter of material treated.

- The compounds I can be converted into the customary formulations, e.g. solutions, emulsions, suspensions, dusts, powders, pastes
40 and granules. The use form depends on the particular purpose; in any case, it should guarantee a fine and uniform distribution of the compound according to the invention.

12

The formulations are prepared in a known manner, e.g. by extending the active ingredient with solvents and/or carriers, if desired using emulsifiers and dispersants, it also being possible to use other organic solvents as auxiliary solvents if water is used as the diluent. Auxiliaries which are suitable are essentially: solvents such as aromatics (e.g. xylene), chlorinated aromatics (e.g. chlorobenzenes), paraffins (e.g. mineral oil fractions), alcohols (e.g. methanol, butanol), ketones (e.g. cyclohexanone), amines (e.g. ethanolamine, dimethylformamide) and water; carriers such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic minerals (e.g. highly-disperse silica, silicates); emulsifiers such as non-ionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers, alkylsulfonates and arylsulfonates) and dispersants such as lignin-sulfite waste liquors and methylcellulose.

Suitable surfactants are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid, phenolsulfonic acid, dibutyl naphthalenesulfonic acid, alkylaryl-sulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates and fatty acids and their alkali metal and alkaline earth metal salts, salts of sulfated fatty alcohol glycol ether, condensates of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensates of naphthalene or of naphthalenesulfonic acid with phenol or formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated isooctylphenol, octylphenol, nonylphenol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ethers, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and methylcellulose.

Substances which are suitable for the preparation of directly sprayable solutions, emulsions, pastes or oil dispersions are mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, e.g. benzene, toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, methanol, ethanol, propanol, butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene, isophorone, strongly polar solvents, e.g. dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and water.

13

Powders, materials for scattering and dusts can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

- 5 Granules, e.g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients to solid carriers. Examples of solid carriers are mineral earths, such as silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, 10 calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.

15

In general, the formulations comprise of from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active ingredient. The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

20

The following are exemplary formulations:

- I. 5 parts by weight of a compound according to the invention are mixed intimately with 95 parts by weight of finely divided kaolin. This gives a dust which comprises 5% by weight 25 of the active ingredient.

- II. 30 parts by weight of a compound according to the invention are mixed intimately with a mixture of 92 parts by weight of pulverulent silica gel and 8 parts by weight of paraffin oil 30 which had been sprayed onto the surface of this silica gel. This gives a formulation of the active ingredient with good adhesion properties (comprises 23% by weight of active ingredient).

35

- III. 10 parts by weight of a compound according to the invention are dissolved in a mixture composed of 90 parts by weight of xylene, 6 parts by weight of the adduct of 8 to 10 mol of ethylene oxide and 1 mol of oleic acid N-monoethanolamide, 2 40 parts by weight of calcium dodecylbenzenesulfonate and 2 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil (comprises 9% by weight of active ingredient).

45

14

- IV. 20 parts by weight of a compound according to the invention are dissolved in a mixture composed of 60 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 5 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 5 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil (comprises 16% by weight of active ingredient).
- 5
- V. 80 parts by weight of a compound according to the invention are mixed thoroughly with 3 parts by weight of sodium diisobutyl-naphthalene- α -sulfonate, 10 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 7 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill (comprises 80% by weight of active ingredient).
- 10
- 15
- VI. 90 parts by weight of a compound according to the invention are mixed with 10 parts by weight of N-methyl- α -pyrrolidone, which gives a solution which is suitable for use in the form of microdrops (comprises 90% by weight of active ingredient).
- 20
- VII. 20 parts by weight of a compound according to the invention are dissolved in a mixture composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.
- 25
- 30
- VIII. 20 parts by weight of a compound according to the invention are mixed thoroughly with 3 parts by weight of sodium diisobutyl-naphthalene- α -sulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill. Finely distributing the mixture in 20,000 parts by weight of water gives a spray mixture which comprises 0.1% by weight of the active ingredient.
- 35
- 40

The active ingredients can be used as such, in the form of their formulations or the use forms prepared therefrom, e.g. in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading, or granules, by means of spraying, atomizing, du-

45

	No.	R ¹	R ²	X	phys. data (m.p. [°C], IR [cm ⁻¹])
5	I-7	CH(CH ₃) ₂	CH ₃	CN	
	I-8	(±) CH(CH ₃)-CH ₂ CH ₃	H	CN	
	I-9	(S) CH(CH ₃)-CH ₂ CH ₃	H	CN	
	I-10	(R) CH(CH ₃)-CH ₂ CH ₃	H	CN	
10	I-11	(±) CH(CH ₃)-CH(CH ₃) ₂	H	CN	
	I-12	(S) CH(CH ₃)-CH(CH ₃) ₂	H	CN	
	I-13	(R) CH(CH ₃)-CH(CH ₃) ₂	H	CN	
	I-14	(±) CH(CH ₃)-CH(CH ₃) ₃	H	CN	
15	I-15	(S) CH(CH ₃)-CH(CH ₃) ₃	H	CN	
	I-16	(R) CH(CH ₃)-CH(CH ₃) ₃	H	CN	
	I-17	(±) CH(CH ₃)-CF ₃	H	CN	
	I-18	(S) CH(CH ₃)-CF ₃	H	CN	
20	I-19	(R) CH(CH ₃)-CF ₃	H	CN	
	I-20	CH ₂ CF ₃	H	CN	
	I-21	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃	OCH ₃	
	I-22	CH(CH ₃) ₂	H	OCH ₃	
25	I-23	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		OCH ₃	182
	I-24	cyclopentyl	H	OCH ₃	
	I-25	CH ₂ CH ₃	CH ₂ CH ₃	OCH ₃	
	I-26	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	OCH ₃	
30	I-27	CH(CH ₃) ₂	CH ₃	OCH ₃	
	I-28	(±) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₃	
	I-29	(S) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₃	
	I-30	(R) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₃	
35	I-31	(±) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₃	
	I-32	(S) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₃	
	I-33	(R) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₃	
	I-34	(±) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₃	
40	I-35	(S) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₃	
	I-36	(R) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₃	
	I-37	(±) CH(CH ₃)-CF ₃	H	OCH ₃	
	I-38	(S) CH(CH ₃)-CF ₃	H	OCH ₃	
45	I-39	(R) CH(CH ₃)-CF ₃	H	OCH ₃	
	I-40	CH ₂ CF ₃	H	OCH ₃	
	I-41	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃	OCH ₂ CH ₃	
	I-42	CH(CH ₃) ₂	H	OCH ₂ CH ₃	
	I-43	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		OCH ₂ CH ₃	135
	I-44	cyclopentyl	H	OCH ₂ CH ₃	

20


	No.	R ¹	R ²	X	phys. data (m.p. [°C], IR [cm ⁻¹])
5	I-45	CH ₂ CH ₃	CH ₂ CH ₃	OCH ₂ CH ₃	
	I-46	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	OCH ₂ CH ₃	
	I-47	CH(CH ₃) ₂	CH ₃	OCH ₂ CH ₃	
	I-48	(±) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₂ CH ₃	
10	I-49	(S) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₂ CH ₃	
	I-50	(R) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₂ CH ₃	
	I-51	(±) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₂ CH ₃	
	I-52	(S) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₂ CH ₃	
15	I-53	(R) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₂ CH ₃	
	I-54	(±) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₂ CH ₃	
	I-55	(S) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₂ CH ₃	
	I-56	(R) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₂ CH ₃	
20	I-57	(±) CH(CH ₃)-CF ₃	H	OCH ₂ CH ₃	
	I-58	(S) CH(CH ₃)-CF ₃	H	OCH ₂ CH ₃	
	I-59	(R) CH(CH ₃)-CF ₃	H	OCH ₂ CH ₃	
	I-60	CH ₂ CF ₃	H	OCH ₂ CH ₃	
25	I-61	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃	OCH(CH ₃) ₂	
	I-62	CH(CH ₃) ₂	H	OCH(CH ₃) ₂	
	I-63	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		OCH(CH ₃) ₂	156
	I-64	cyclopentyl	H	OCH(CH ₃) ₂	
30	I-65	CH ₂ CH ₃	CH ₂ CH ₃	OCH(CH ₃) ₂	102
	I-66	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	OCH(CH ₃) ₂	
	I-67	CH(CH ₃) ₂	CH ₃	OCH(CH ₃) ₂	
	I-68	(±) CH(CH ₃)-CH ₂ CH ₃	H	OCH(CH ₃) ₂	
35	I-69	(S) CH(CH ₃)-CH ₂ CH ₃	H	OCH(CH ₃) ₂	
	I-70	(R) CH(CH ₃)-CH ₂ CH ₃	H	OCH(CH ₃) ₂	
	I-71	(±) CH(CH ₃)-CH(CH ₃) ₂	H	OCH(CH ₃) ₂	
	I-72	(S) CH(CH ₃)-CH(CH ₃) ₂	H	OCH(CH ₃) ₂	
40	I-73	(R) CH(CH ₃)-CH(CH ₃) ₂	H	OCH(CH ₃) ₂	
	I-74	(±) CH(CH ₃)-CH(CH ₃) ₃	H	OCH(CH ₃) ₂	
	I-75	(S) CH(CH ₃)-CH(CH ₃) ₃	H	OCH(CH ₃) ₂	
	I-76	(R) CH(CH ₃)-CH(CH ₃) ₃	H	OCH(CH ₃) ₂	
45	I-77	(±) CH(CH ₃)-CF ₃	H	OCH(CH ₃) ₂	
	I-78	(S) CH(CH ₃)-CF ₃	H	OCH(CH ₃) ₂	
	I-79	(R) CH(CH ₃)-CF ₃	H	OCH(CH ₃) ₂	
	I-80	CH ₂ CF ₃	H	OCH(CH ₃) ₂	
	I-81	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃	OCH ₂ CH=CH ₂	
	I-82	CH(CH ₃) ₂	H	OCH ₂ CH=CH ₂	

No.	R ¹	R ²	X	phys. data (m.p. [°C], IR[cm ⁻¹])
5	I-83	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		2990, 1655, 985
	I-84	cyclopentyl	H	OCH ₂ CH=CH ₂
	I-85	CH ₂ CH ₃	CH ₂ CH ₃	OCH ₂ CH=CH ₂
	I-86	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	OCH ₂ CH=CH ₂
10	I-87	CH(CH ₃) ₂	CH ₃	OCH ₂ CH=CH ₂
	I-88	(±) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₂ CH=CH ₂
	I-89	(S) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₂ CH=CH ₂
	I-90	(R) CH(CH ₃)-CH ₂ CH ₃	H	OCH ₂ CH=CH ₂
15	I-91	(±) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₂ CH=CH ₂
	I-92	(S) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₂ CH=CH ₂
	I-93	(R) CH(CH ₃)-CH(CH ₃) ₂	H	OCH ₂ CH=CH ₂
	I-94	(±) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₂ CH=CH ₂
20	I-95	(S) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₂ CH=CH ₂
	I-96	(R) CH(CH ₃)-CH(CH ₃) ₃	H	OCH ₂ CH=CH ₂
	I-97	(±) CH(CH ₃)-CF ₃	H	OCH ₂ CH=CH ₂
	I-98	(S) CH(CH ₃)-CF ₃	H	OCH ₂ CH=CH ₂
25	I-99	(R) CH(CH ₃)-CF ₃	H	OCH ₂ CH=CH ₂
	I-100	CH ₂ CF ₃	H	OCH ₂ CH=CH ₂
	I-101	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃	OCHF ₂
	I-102	CH(CH ₃) ₂	H	OCHF ₂
30	I-103	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		138
	I-104	cyclopentyl	H	OCHF ₂
	I-105	CH ₂ CH ₃	CH ₂ CH ₃	OCHF ₂
	I-106	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	OCHF ₂
35	I-107	CH(CH ₃) ₂	CH ₃	OCHF ₂
	I-108	(±) CH(CH ₃)-CH ₂ CH ₃	H	OCHF ₂
	I-109	(S) CH(CH ₃)-CH ₂ CH ₃	H	OCHF ₂
	I-110	(R) CH(CH ₃)-CH ₂ CH ₃	H	OCHF ₂
40	I-111	(±) CH(CH ₃)-CH(CH ₃) ₂	H	OCHF ₂
	I-112	(S) CH(CH ₃)-CH(CH ₃) ₂	H	OCHF ₂
	I-113	(R) CH(CH ₃)-CH(CH ₃) ₂	H	OCHF ₂
	I-114	(±) CH(CH ₃)-CH(CH ₃) ₃	H	OCHF ₂
45	I-115	(S) CH(CH ₃)-CH(CH ₃) ₃	H	OCHF ₂
	I-116	(R) CH(CH ₃)-CH(CH ₃) ₃	H	OCHF ₂
	I-117	(±) CH(CH ₃)-CF ₃	H	OCHF ₂
	I-118	(S) CH(CH ₃)-CF ₃	H	OCHF ₂

22

No.	R ¹	R ²	X	phys. data (m.p. [°C], IR [cm ⁻¹])
I-119	(R) CH(CH ₃)-CF ₃	H	OCHF ₂	
I-120	CH ₂ CF ₃	H	OCHF ₂	

Table II



II

No.	R ¹	R ²	phys. data (m.p. [°C])
II-1	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃	136
II-2	CH(CH ₃) ₂	H	143
II-3	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		150
II-4	cyclopentyl	H	62
II-5	CH ₂ CH ₃	CH ₂ CH ₃	167
II-6	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	148
II-7	CH(CH ₃) ₂	CH ₃	151
II-8	(±) CH(CH ₃)-CH ₂ CH ₃	H	150
II-9	(S) CH(CH ₃)-CH ₂ CH ₃	H	147
II-10	(R) CH(CH ₃)-CH ₂ CH ₃	H	147
II-11	(±) CH(CH ₃)-CH(CH ₃) ₂	H	145
II-12	(S) CH(CH ₃)-CH(CH ₃) ₂	H	145
II-13	(R) CH(CH ₃)-CH(CH ₃) ₂	H	145
II-14	(±) CH(CH ₃)-CH(CH ₃) ₃	H	188
II-15	(S) CH(CH ₃)-CH(CH ₃) ₃	H	191
II-16	(R) CH(CH ₃)-CH(CH ₃) ₃	H	191
II-17	(±) CH(CH ₃)-CF ₃	H	179
II-18	(S) CH(CH ₃)-CF ₃	H	172
II-19	(R) CH(CH ₃)-CF ₃	H	172
II-20	CH ₂ CF ₃	H	211
II-21	H	H	250

Examples of the action against harmful fungi.

The fungicidal action of the compounds of the formula I was demonstrated by the following experiments:

23

The active compounds, separately or together, were formulated as a 10% emulsion in a mixture of 70% by weight of cyclohexanone, 20% by weight of Nekanil® LN (Lutensol® AP6, wetting agent having emulsifying and dispersant action based on ethoxylated alkylphenols) and 10% by weight of Wettol® EM (nonionic emulsifier based on ethoxylated castor oil) and diluted with water to the desired concentration.

Use example 1: Action on *Alternaria solani* in tomatoes

10 Leaves of pot grown tomato seedlings of the "Große Fleischtomate St. Pierre" variety were sprayed with aqueous liquors made from a stock solution consisting of 10% of active ingredient, 63% of cyclohexanone, and 27% of emulsifier. After 24 hours the leaves
15 were infected with a zoospore suspension of *Alternaria solani* (1.7×10^6 spores per ml of a 2% strength biomalt solution). The plants were then placed in a water vapour-saturated chamber. After 5 days the disease had spread to such a great extent on the untreated plants that the fungicidal activity of the substances
20 could be assessed.

In this test, the plants which had been treated with 63 ppm of compounds I-1, I-3, and I-23 showed an infection of up to 3%, whereas the untreated plants were infected to 100%.


25 Use Example 2: Action on *Plasmopara viticola*

Leaves of potted vines of the "Müller Thurgau" variety were sprayed with aqueous liquors made from a stock solution consisting of 10% of active ingredient, 63% of cyclohexanone, and 27%
30 of emulsifier. To assess the duration of action, the plants were set up, after the sprayed-on layer had dried, in the greenhouse for 8 days. Then the leaves were infected with a zoospore suspension of the fungus *Plasmopara viticola*, first placed in a vapour-saturated chamber at 24°C, and then kept for 5 days in a greenhouse at 20 to 30°C. To accelerate and intensify the sporangio-
35 phore discharge, the plants were then again placed in the moist chamber for 16 hours. The extent of fungus attack was then assessed on the undersides of the leaves.

40 In this test, the plants which had been treated with 250 ppm of compounds I-1 and I-2 showed an infection of up to 20%, whereas the untreated plants were infected to 85%.

45 Active compounds A to E known from EP-A 550 113, WO-A 99/41255, and US 5,994,360 resp., were used as comparison compounds in the following comparison tests:

24



5

No.	known from	R	X
A	EP-A 550 113, No. 106	isopropylamino	chloro
B	EP-A 550 113, No. 111	diethylamino	chloro
10 C	WO-A 99/41255, No. 38	cyclohexyl	methoxy
D	WO-A 99/41255, No. 39	cyclohexyl	cyano
E	US 5,994,360, Ex. 1B	4-CH ₃ -piperidine	methyl

15 Comparison test 1: Fungicidal control of early blight on tomatoes (*Alternaria solani*)

Young seedlings of tomato plants of the variety "Große Fleischtomate St. Pierre" were grown in pots to the 2 to 4 leaf stage.

20 These plants were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient mentioned in the table below, prepared from a stock solution containing 10 % of the active ingredient, 85 % cyclohexanone and 5 % emulsifier. The next day, the treated plants were inoculated with an aqueous

25 suspension of *Alternaria solani* containing 0.2×10^6 spores per ml. Then the trial plants were immediately transferred to a humid chamber. After 6 days at 20 to 23°C and a relative humidity close to 100 %, the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

30 In this trial, the plants which have been treated with 250 ppm of compounds I-3 and I-23, resp., showed an infection of not more than 7 %, whereas the plants treated with 250 ppm of comparison compounds C and D, resp., were infected to at least 60 %, and the

35 untreated plants were infected to 100 %.

In another trial, the plants which have been treated with 16 ppm of compound I-3 showed an infection of 15 %, whereas the plants treated with 16 ppm of comparison compound E, were infected to

40 30 %, and the untreated plants were infected to 90 %.

Comparison test 2: Control of net blotch on barley caused by *Pyrenophora teres*

45 The first fully developed leaves of pot grown barley of the variety "Igri" were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient or their mix-

25

ture mentioned in the table below, prepared from a stock solution containing 10 % of the active ingredient, 85 % cyclohexanone and 5 % emulsifier. The next day the treated plants were inoculated with an aqueous spore suspension of *Pyrenophora teres tritici* containing 0.02×10^6 spores/ml. Then the trial plants were immediately transferred to a humid chamber in the greenhouse. After 6 days of cultivation at 20 to 24°C and a relative humidity close to 100 %, the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

10

In this trial, the plants which have been treated with 250 ppm of compounds I-3 and I-23, resp., showed an infection of not more than 7 %, whereas the plants treated with 250 ppm of comparison compounds C and D, resp., and the untreated plants were infected to 100 %.

15

Comparison trial 3: Fungicidal control of leaf blotch on wheat caused by *Septoria tritici* (protective)

20 Leaves of pot-grown wheat seedling of the variety "Riband" were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient as described below prepared from a stock solution containing 10 % of the active ingredient, 85 % cyclohexanone and 5 % emulsifier. The plants were allowed to air-dry. At the following day the plants were inoculated with an aqueous spore suspension of *Septoria tritici* containing 2.0×10^6 spores/ml. Then the trial plants were immediately transferred to a humid chamber. After 2 weeks at 18 to 22°C and a relative humidity close to 100 % the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

30

In this trial, the plants which have been treated with 63 ppm of compounds I-3, I-5 and I-23, resp., showed no infection, whereas the plants treated with 63 ppm of comparison compounds B, C and D, resp., were infected from 20 to 60 %, and the untreated plants were infected to 90 %.

35

Comparison test 4: Fungicidal control of rice blast caused by *Pyricularia oryzae* (protective)

40

Leaves of pot-grown rice seedling of the variety "Tai-Nong 67" were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient as described below prepared from a stock solution containing 10 % of the active ingredient, 85 % cyclohexanone and 5 % emulsifier. The plants were allowed to air-dry. At the following day the plants were inoculated with an aqueous spore suspension of *Pyricularia oryzae* containing

45

26

1.0 x 10⁶ spores/ml. Then the trial plants were immediately transferred to a humid chamber. After 6 days at 22 to 24°C and a relative humidity close to 100 % the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

5

In this trial, the plants which have been treated with 250 ppm of compounds I-3, I-5, I-23, and I-63 resp., showed an infection of up to 30 %, whereas the plants treated with 250 ppm of comparison compounds A, B, and E, resp., were infected from 60 to 80 %, and
10 the untreated plants were infected to 90 to 100 %.

Comparison test 5: Protective control of powdery mildew on cucumber

15 Cucumber seedlings of the variety "Chinesische Schlange" were grown in pots to the 2 leaf stage. The plants were then sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient as described below, prepared from a stock solution containing 10 % of the active ingredient, 85 % cy-
20 clohexanone and 5 % emulsifier. The next day the treated plants were inoculated with an aqueous spore suspension of cucumber powdery mildew (*Sphaerotheca fuliginea*). Then the trial plants were cultivated in a greenhouse at temperatures between 20 and 24°C and a relative humidity between 60 and 80 %. After 8 days the extent
25 of fungal attack on the leaves was visually assessed as % diseased leaf area.

In this trial, the plants which have been treated with 16 ppm of compound I-23 showed no infection, whereas the plants treated
30 with 16 ppm of comparison compound E were infected to 90 %, and the untreated plants were infected to 100 %.


35

40

45

Claims:

1. 6-(2-Chloro-6-fluoro-phenyl)-triazolopyrimidines of formula I



I

in which

R¹ and R² independently denote hydrogen or

C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₄-C₁₀-alkadienyl, or C₁-C₁₀-haloalkyl,

C₃-C₈-cycloalkyl, C₅-C₁₀-bicycloalkyl, phenyl, naphthyl, or

5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or

5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or

where R¹ and R² radicals may be unsubstituted or partially or fully halogenated or may carry one to three groups R^a,

R^a is halogen, cyano, nitro, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, di-C₁-C₆-alkylamino, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₂-C₆-alkynyl, C₃-C₆-alkynyloxy and C₁-C₄-alkylenedioxy, which may be halogenated; or

R¹ and R² together with the interjacent nitrogen atom represent a 5- or 6-membered heterocyclic ring, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, which may be substituted by one to three R^a radicals;

X is cyano, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy or C₃-C₈-alkenyloxy.

28

2. Compounds of formula I according to claim 1, in which

R¹ is straight chained or branched C₁-C₆-alkyl, C₁-C₆-halo-alkyl, straight chained or branched C₂-C₆-alkenyl,
 5 C₃-C₆-cycloalkyl, or C₅-C₉-bicycloalkyl, and

R² is hydrogen or C₁-C₆-alkyl, or

10 R¹ and R² together with the interjacent nitrogen atom represent a heterocyclic ring with 5 or 6 carbon atoms being optionally substituted with one or two C₁-C₆-alkyl groups.

3. Compounds according to claims 1 or 2 in which R² is hydrogen.


15

4. Compounds according to claims 1 to 3 in which X is methoxy.

5. Compounds according to claims 1 to 3 in which X is cyano.

- 20 6. A process for the preparation of compounds of formula I as defined in claims 1 to 5 which comprises reacting 5-halogen-6-phenyl-triazolopyrimidines of formula II

25



II

in which Y is halogen with a compound of formula III

30

M-X

III

in which M is an ammonium-, tetraalkylammonium-, alkalimetal- or earth metal cation and X is as defined in formula I to
 35 produce compounds of formula I.

7. The use of compounds of formula II as defined in claim 6 as intermediate.

- 40 8. A composition suitable for controlling phytopathogenic fungi, comprising a solid or liquid carrier and a compound of the formula I as claimed in claim 1.

45

29

9. A method for controlling phytopathogenic fungi, which comprises treating the fungi or the materials, plants, the soil or the seed to be protected against fungal attack with an effective amount of a compound of the formula I as claimed in claim 1.

10

15

20

25

30

35

40

45

INTERNATIONAL SEARCH REPORT

Inventor's Application No.
EPO/EP 02/03830

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D487/04 A01N43/90 //(C07D487/04,249:00,239:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 41255 A (AMERICAN CYANAMID CO) 19 August 1999 (1999-08-19) page 2 page 5 examples 38,39,52,54; table 1	1
Y	US 5 994 360 A (PFRENGLE WALDEMAR) 30 November 1999 (1999-11-30) cited in the application column 12 -column 13 examples 4-6,9,10,12,15,18; table 1	1
Y	FR 2 784 991 A (AMERICAN CYANAMID CO) 28 April 2000 (2000-04-28) page 3, line 25 -page 6, line 25 page 6, line 4 - line 5	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

20 June 2002

Date of mailing of the international search report

28/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Goss, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/03830

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9941255	A	19-08-1999	US 6020338 A	01-02-2000
			AU 2595299 A	30-08-1999
			BR 9907863 A	24-10-2000
			CA 2320304 A1	19-08-1999
			CN 1292790 T	25-04-2001
			CZ 20002933 A3	17-04-2002
			EP 1054888 A1	29-11-2000
			HU 0100885 A2	28-06-2001
			JP 2002503664 T	05-02-2002
			PL 342576 A1	18-06-2001
			WO 9941255 A1	19-08-1999
US 5994360	A	30-11-1999	NONE	
FR 2784991	A	28-04-2000	FR 2784991 A1	28-04-2000